


SUPREP BOWEL PREPARATION

Digestive Health Institute



SUPREP BOWEL PREPARATION

Your preparation drink was sent to your pharmacy. Please ensure you pick it up within 7 days of being notified from your pharmacy that it is ready.

DO NOT WAIT TO PICK IT UP CLOSER TO YOUR SCHEDULED DATE

It is important that you follow the preparation instructions listed here. Disregard any other instructions from the preparation packaging. Your procedure may be canceled if instructions here are not followed.


Please make arrangements for a responsible adult to accompany you after completion of the procedure. Your procedure may be subject to cancellation if you have no transportation. Taxi and rideshare options are not acceptable unless you have an accompanying responsible adult.

You should expect to be at the procedure location for approximately 3 hours from time of registration to discharge.

If you have further questions, please call your Nuvance Health Gastroenterology practice at:

Danbury: (203)739-7038
Norwalk: (203) 852-2278

For questions regarding insurance coverage for your procedure, please contact your insurance company directly, or our Financial Clearance Department at 203-794-5350.



MEDICATION MANAGEMENT

If you take any medications that increase your risk of bleeding and were not told when to stop the medication by your gastroenterologist, please check with the prescribing doctor prior to the procedure.

EXAMPLES:

BRILINTA (TICAGRELOR)
ELIQUIS (APIXABAN)
LIXIANA (EDOXABAN)
LOVENOX (ENOXAPARIN)
XARELTO (RIVAROXABAN)

PLAVIX (CLOPIDOGREL)
PRADAXA (DABIGATRAN)
PRASUGREL (EFFIENT)
WARFARIN (COUMADIN)

Please note: OK to continue 81mg aspirin (baby aspirin).

For patients with diabetes, metabolic diseases, or are taking a weight loss medication:

- If you take insulin (injectable or pump), and were not given instructions by the gastroenterologist, please see your prescribing physician for instructions on dosage adjustment.
- If you take oral diabetes medications (except GLP-1 agonist/weight loss medications-see below), please do not take them the morning of your procedure. You may resume them after your procedure with a regular diet.
- For patients taking GLP-1 agonists (often called weight loss medications):

Drug	Directions of Stopping Drug	RESUMING DRUG AFTER PROCEDURE
Once weekly GLP-1 (injection) Semaglutide (Ozempic, Wegovy) Dulaglutide (Trulicity) Tirzapatide (Mounjaro, Zepbound) Exenatide (Bydureon)	Hold for a minimum of 8 days prior to procedure (including day of procedure)	Resume next day or next scheduled dose
Once daily oral GLP-1 Semaglutide (Rybelsus)	Do not take on morning of procedure	Resume next scheduled dose
Once daily GLP-1 (injection) Liraglutide (Victoza, Saxenda)	Do not take on morning of procedure	Resume next scheduled dose
Twice daily GLP-1 (injection) Exenatide (Byetta)	Do not take night before, or on morning of procedure	Resume next scheduled dose



5 DAYS PRIOR TO COLONOSCOPY


- Stop taking iron supplements, herbal medications and all vitamins unless directed by your gastroenterologist.

3 DAYS PRIOR TO COLONOSCOPY

- Avoid eating high residue foods. These includes all raw fruits, raw vegetables, seeds, nuts, quinoa, wheat bran, lentils, corn, vegetables with seeds, poppy seeds, sesame seeds, pumpkin seeds, etc.
 - Cooked and steamed vegetables are OK!
- Confirm your transportation.
- Confirm the location of your procedure and secure appropriate directions and time of arrival.

1 DAY PRIOR TO COLONOSCOPY

- DO NOT eat any solid food
- DO NOT drink alcohol
- Drink plenty of clear liquids to ensure adequate hydration. **DO NOT DRINK RED OR PURPLE LIQUIDS (ALL OTHER COLORS ARE OK)**
- Examples of clear liquids include:
 - Water
 - Clear soup, broth, bouillon or consommé
 - Sports drinks
 - Gelatin such as Jell-O, flavored ices
 - Clear juice (apple, lemonade, white grape juice, white cranberry juice)
 - Tea or coffee (black)
 - Ginger-ale, lemon-lime soda
 - Popsicles
- **At 6pm the night before procedure:** Pour one bottle of Suprep into the clear container from the Suprep box. Fill the container to the top with water and drink the mixture.
- 30 minutes after completing the mixture, fill same clear container with water (or another clear liquid of your choice- see above) twice and drink slowly over the course of 1 hour to ensure proper flushing of the preparation.



DAY OF THE COLONOSCOPY

- **5 hours prior to scheduled arrival time:**
 - Repeat exact same procedure as the night before: mix the second bottle of Suprep with water and drink. Then slowly drink 2 containers of water (or another clear liquid of your choice- see above) over the next hour.
- All prep and clear liquids should be completed 2 hours before the scheduled arrival time
- No chewing gum or hard candy the morning of the procedure
- Can take heart and blood pressure medications with a small sip of water up to 2 hours before your arrival time, or after your procedure is completed
- **Do not take oral diabetic medications or diuretic (water pill) medications**
- Bring your current list of medications
- Bring your defibrillator/pacemaker card if you have one
- **DO NOT bring valuables or jewelry**
- **DO NOT wear contact lenses**
- Wear comfortable clothing

You should expect to be at the procedure location for approximately 3 hours from time of registration to discharge.

If you have further questions, please call your Nuvance Health Gastroenterology practice at:

Danbury: (203)739-7038
Norwalk: (203) 852-2278